arXiv:1506.08980v1 [math.GN] 30 Jun 2015

FUNCTIONALLY COUNTABLE SUBALGEBRAS AND SOME PROPERTIES OF BANASCHEWSKI COMPACTIFICATION

ALIREZA OLFATI

Abstract. Let X be a zero-dimensional space and $C_c(X)$ be the set of all continuous real valued functions on X with countable image. In this article we denote by $C_c^K(X)$ (resp., $C_c^\psi(X)$) the set of all functions in $C_c(X)$ with compact (resp., pseudocompact) support. First, we observe that $C_c^K(X) = O_c^{\beta_0 X\setminus X}$ (resp., $C_c^\psi(X) = M_c^{\beta_0 X\setminus \upsilon_0 X}$). This implies that for an $\mathbb{N}$-compact space X, the intersection of all free maximal ideals in $C_c(X)$ equals to $C_c^K(X)$, i.e., $M_c^{\beta_0 X\setminus X} = C_c^K(X)$. Afterwards, by applying methods of functionally countable subalgebras, we observe some results in remainder of Banaschewski compactification. It is shown that for a zero-dimensional non pseudocompact space X, the set $\beta_0 X\setminus \upsilon_0 X$ has cardinality at least $2^{2^{\aleph_0}}$. Moreover, for a locally compact and $\mathbb{N}$-compact space X, the remainder $\beta_0 X\setminus X$ is an almost P-space. These results leads us to find a class of Parovičenko spaces in Banaschewski compactification of a non pseudocompact zero-dimensional space. We conclude with a theorem which gives a lower bound for the cellularity of subspaces $\beta_0 X\setminus \upsilon_0 X$ and $\beta_0 X\setminus X$, whenever X is a zero-dimensional, locally compact space which is not pseudocompact.

1. Preliminaries

We recall that a zero-dimensional topological space is a Hausdorff space with a base consisting of clopen sets. Mrowka showed that X is zero-dimensional if and only if it can be embedded into the product space $\mathbb{N}^\kappa$, where $\mathbb{N}$ is the set of natural numbers with discrete topology and κ an arbitrary cardinal number.

We also recall that a topological space X is $\mathbb{N}$-compact if it can be embedded as a closed subset of the product space $\mathbb{N}^\kappa$, for some cardinal number κ, see [3, 13, 14, 15, 16, 17] for more details on this subject.

For every zero-dimensional space X, there exists an $\mathbb{N}$-compact space $\upsilon_0 X$ such that X is dense in it and every continuous function $f : X \to Y$, with Y an $\mathbb{N}$-compact space, has a unique extension $f^* : \upsilon_0 X \to Y$. We can replace an arbitrary $\mathbb{N}$-compact space Y by the fixed discrete space $\mathbb{Z}$ (the set of integer numbers) and have the following characterization of $\mathbb{N}$-compactification of a zero-dimensional space, see [19, 5.4 (d)].

Theorem 1.1. An $\mathbb{N}$-compact extension T of a zero-dimensional space X is homeomorphic to $\upsilon_0 X$ if and only if for each continuous function $f : X \to \mathbb{Z}$, there exists $F : T \to \mathbb{Z}$ such that $F|_X = f$.

2000 *Mathematics Subject Classification.* Primary 54C30; 54C40; 54A25; Secondary 54D40, 54D60, 54G05.

Key words and phrases. Zero-dimensional space, Strongly zero-dimensional Space, $\mathbb{N}$-compact space, Banaschewski compactification, Pseudocompact space, Functionally countable subalgebra, Support, Cellularity, Remainder, Almost P-space, Parovičenko space.

We remind that a Tychonoff space X is strongly zero-dimensional if and only if every two disjoint zero-sets in X are separated by a clopen partition. It is well known that every strongly zero-dimensional real compact space is $\mathbb{N}$-compact. Since every countable subset of $\mathbb{R}$ (the set of real numbers) is Lindelöff and zero-dimensional, it should be strongly zero-dimensional. This fact implies that every countable subset of $\mathbb{R}$ is $\mathbb{N}$-compact. So we have the following lemma.

Lemma 1.2. Let X be a zero-dimensional Hausdorff space. For each continuous function $f : X \to \mathbb{R}$ with countable image, there exists an extension $f^* : \upsilon_0 X \to \mathbb{R}$ such that the image of f^* equals to the image of f.

For an arbitrary Tychonoff space X, we denote by $C_c(X)$ the set of all continuous real valued functions on X with countable image. The set $C_c(X)$ forms a subring with pointwise addition and multiplication. Ghadermazi, Karamzadeh and Namdari showed in [5] that for a Tychonoff space X there exists a zero-dimensional space Y such that $C_c(X) \cong C_c(Y)$ as a ring isomorphism. Motivated by this fact, in the present article we restrict our attension to zero-dimensional spaces. The reader could find all prerequisites and unfamiliar notions for this subring in [5].

For a zero-dimensional space X, let $\beta_0 X$ denote its Banaschewski compactification. We recall that $\beta_0 X$ is the unique (up to homeomorphism) zero-dimensional compact space containing X as a dense subset such that every continuous two-valued function $f : X \to \{0, 1\}$ has an extension to $\beta_0 X$, see [19, 4.7, Corollary (f)]. It was shown that X is strongly zero-dimensional if and only if $\beta_0 X$ is homeomorphic to βX, i.e., βX is zero-dimensional, see [2].

Dowker gave an example of a zero-dimensional space X in which βX is not zero-dimensional and hence $\beta X \neq \beta_0 X$, see [19, exercise 4V]. The structure of $\beta_0 X$ is related to the clopen ultrafilters defined on X. Indeed $\beta_0 X$ is homeomorphic to the set of all clopen ultrafilters equipped with the stone topology, see [19, 4.7]. In [19, exercise 5E], it was also given an outline for recovering $\upsilon_0 X$ as a subspace of all clopen ultrafilters on X which have countable intersection property. Therefore we have $X \subseteq \upsilon_0 X \subseteq \beta_0 X$. Note that for $p \in \beta_0 X \setminus \upsilon_0 X$, there exists a sequence $\{V_n : n \in \mathbb{N}\}$ of clopen neighborhoods of p in $\beta_0 X$ such that $\cap_{n=1}^{\infty} V_n$ does not meet X.

This article consists of two parts. In part one, by applying previous notions, we characterize some important ideals in $C_c(X)$. It is shown that the set of all functions in $C_c(X)$ with compact (resp., pseudocompact) support equals to the ideal $O_c^{\beta_0 X \setminus X}$ (resp., $M_c^{\beta_0 X \setminus \upsilon_0 X}$). It is shown that for an $\mathbb{N}$-compact space X, the intersection of all free maximal ideals in $C_c(X)$ equals to the ideal $C_c^K(X)$. In part two, we apply the subring $C_c(X)$ and some related notions to find some information about $\beta_0 X$. For example, we will give the least cardinality of the remainder $\beta_0 X \setminus \upsilon_0 X$, i.e., the set of all clopen ultrafilters on X which do not have countable intersection property. It is shown that for a non pseudocompact space X, there are at least $2^{2^{\aleph_0}}$ such clopen ultrafilters on X. We shall show that if X is locally compact and $\mathbb{N}$-compact, then $\beta_0 X \setminus X$ is an almost P-space, i.e., a space for which the interior of every zero-set is nonempty, see [10] and [20]. We show that zero-sets of $\beta_0 X$ which do not meet X are Parovičenko spaces. In the end, it is shown that whenever X is zero-dimensional, locally compact which is not pseudocompact, the cellularity of the spaces $\beta_0 X \setminus \upsilon_0 X$ and $\beta_0 X \setminus X$ are at least $2^{\aleph_0}$. We close this section with the following two results which are useful in the rest of this article.

Lemma 1.3. (a) For a sequence $\{U_n : n \in \mathbb{N}\}$ of clopen subsets of X, there exists $f \in C_c(X)$ such that $Z(f) = \cap_{n=1}^{\infty} U_n$.
(b) If $f \in C_c(X)$, then there exists a sequence $\{W_n : n \in \mathbb{N}\}$ of clopen subsets of X such that $Z(f) = \cap_{n=1}^{\infty} W_n$.

Proof. (a) With out lose of generality, we could assume that $U_1 \supseteq U_2 \supseteq U_3 \cdots$ be a decreasing sequence of clopen sets of X. Now define

$$f = \sum_{n=1}^{\infty} \frac{1}{2^n} \chi_{(X \setminus U_n)},$$

where $\chi_{X \setminus U_n}$ is the characteristic function of the clopen set $X \setminus U_n$. It is easy to see that $f(X) \subseteq \{0\} \cup \{\frac{1}{2^n} : n \in \mathbb{N} \cup \{0\}\}$. Therefore $f \in C_c(X)$ and $Z(f) = \cap_{n=1}^{\infty} U_n$.
(b) Suppose that $Z \in Z_c[X]$. Consider $0 < f \in C_c(X)$ such that $Z = Z(f)$. Choose a decreasing sequence $r_1 > r_2 > \cdots > r_n > \cdots$ of real numbers which tends to zero and for each $n \in \mathbb{N}$, $r_n \notin f(X)$. For each $n \in \mathbb{N}$, define $W_n = f^{-1}[0, r_n)$. Then for each $n \in \mathbb{N}$, W_n is clopen in X and $Z(f) = \cap_{n=1}^{\infty} W_n$. So we are done. □

Corollary 1.4. Let X be a zero-dimensional space. For $f \in C_c(X)$ there exists $F \in C_c(\beta_0 X)$ such that $Z(f) = Z(F) \cap X$.

Proof. By part (b) of Lemma 1.3, there exists a sequence $\{W_n : n \in \mathbb{N}\}$ of clopen subsets of X such that $Z(f) = \cap_{n=1}^{\infty} W_n$. For each $n \in \mathbb{N}$, $cl_{\beta_0 X} W_n$ is a clopen subset in $\beta_0 X$. Part (a) of Lemma 1.3 implies that there exists $F \in C_c(X)$ such that $Z(F) = \bigcap_{n=1}^{\infty} cl_{\beta_0 X} W_n$. Clearly $Z(f) = Z(F) \cap X$. □

2. Characterisation of some special ideals in $C_c(X)$.

In the begining of this section we characterize maximal ideals of $C_c(X)$. This characterization leads us to specify the ideal which consists of all functions in $C_c(X)$ with compact (resp., pseudocompact) support. We remind that in [1] it is shown that the space of maximal ideals of $C_c(X)$ is isomorphic to $\beta_0 X$. But we need to settle the internal characterization of maximal ideals of $C_c(X)$. In the following we could observe the counterpart of Gelfand-Kolmogoroff theorem in rings of continuous functions. First, we need a lemma which is essential for characterizing all maximal ideals of this subring.

Lemma 2.1. Let X be zero-dimensional. For each two $f, g \in C_c(X)$ we have

$$cl_{\beta_0 X} \left(Z(f) \cap Z(g)\right) = cl_{\beta_0 X} Z(f) \cap cl_{\beta_0 X} Z(g).$$

Proof. Case 1. Suppose $Z(f) \cap Z(g) = \emptyset$. There exists $h \in C_c(X)$ such that $Z(f) = Z(h)$ and $Z(g) = h^{-1}(1)$. Choose $0 < r < 1$ such that $r \notin h(X)$ and put $U = h^{-1}(-\infty, r)$. The subset U is clopen in X, $Z(f) \subseteq U$ and $Z(g) \subseteq X \setminus U$. Note that X is two-embedded in $\beta_0 X$ and this implies that $cl_{\beta_0 X} U \cap cl_{\beta_0 X}(X \setminus U) = \emptyset$ and then $cl_{\beta_0 X} Z(f) \cap cl_{\beta_0 X} Z(g) = \emptyset$.
Case 2. Evidently $cl_{\beta_0 X} \left(Z(f) \cap Z(g)\right) \subseteq cl_{\beta_0 X} Z(f) \cap cl_{\beta_0 X} Z(g)$. For $p \in cl_{\beta_0 X} Z(f) \cap cl_{\beta_0 X} Z(g)$, assume that p does not belong to $cl_{\beta_0 X} \left(Z(f) \cap Z(g)\right)$. There exists a clopen set $U \subseteq \beta_0 X$ such that $p \in U$ and $U \cap (cl_{\beta_0 X} \left(Z(f) \cap Z(g)\right)) = \emptyset$. The subset $V = U \cap X$ is clopen in X and also is the zero-set of the characteristic function $\chi_{X \setminus V}$, i.e., $Z(\chi_{X \setminus V}) = V$. Then $V \cap Z(f) \cap Z(g) = \emptyset$. By case 1, we have

$$cl_{\beta_0 X}(V \cap Z(f)) \cap cl_{\beta_0 X}(V \cap Z(g)) = \emptyset,$$

and hence p does not belong to at least one of them, say for example $p \notin cl_{\beta_0 X}(V \cap Z(f))$. Choose a neighborhood W of p such that $W \cap V \cap Z(f) = \emptyset$ and hence $W \cap U \cap Z(f) = \emptyset$. But $W \cap U$ is a neighborhood of p and must intersects $Z(f)$. This is a contradiction and the proof is complete. □

Theorem 2.2. [Gelfand-Kolmogoroff] The Maximal ideals of $C_c(X)$ are in one to one correspondence with the points of $\beta_0 X$ and are given by

$$M_c^p = \{f \in C_c(X) : p \in cl_{\beta_0 X} Z(f)\},$$

for $p \in \beta_0 X$.

Proof. First we show that for each $p \in \beta_0 X$, M_c^p is a maximal ideal. By Lemma 2.1, one can verify that M_c^p forms an ideal. Suppose on the contrary that there exists a maximal ideal M which properly contains M_c^p. Therefore there exists $f \in M$ such that $p \notin cl_{\beta_0 X} Z(f)$. Note that $\beta_0 X$ is zero-dimensional. Hence there exists a clopen subset U in $\beta_0 X$ such that $p \in U$ and $U \cap cl_{\beta_0 X} Z(f) = \emptyset$. The set $V = U \cap X$ is clopen in X and clearly $p \in cl_{\beta_0 X} V$. Since V is the zero-set of the characteristic function $\chi_{X \setminus V} \in C_c(X)$, therefore $\chi_{X \setminus V} \in M_c^p$. Consider the function $g = \chi_{X \setminus V} + f^2$. Obviously $Z(g) = Z(\chi_{X \setminus V}) \cap Z(f) = V \cap Z(f) = \emptyset$. Therefore M contains a unit of $C_c(X)$ which implies that $M = C_c(X)$.
Now we must show that each maximal ideal M in $C_c(X)$ has this form. By Lemma 2.1, the set $\{cl_{\beta_0 X} Z(f) : f \in M\}$ is a family of closed subsets with the finite intersection property and since $\beta_0 X$ is compact, there exists $p \in \bigcap_{f \in M} cl_{\beta_0 X} Z(f)$. Therefore $M \subseteq M_c^p$ and hence $M = M_c^p$. □

Suppose that $p \in \beta_0 X$. We define the set O_c^p as follows.

$$O_c^p = \{f \in C_c(X) : p \in int_{\beta_0 X} cl_{\beta_0 X} Z(f)\}.$$

One could observe that O_c^p forms an ideal of $C_c(X)$ and the only maximal ideal containing O_c^p is M_c^p. For $p \in \beta_0 X$, in the following we characterize the set $Z[O_c^p]$. Note that in general, all bounded functions on X could not be extended to $\beta_0 X$. But the following proposition allows us to extend zero-sets of O_c^p.

Proposition 2.3. For each $p \in \beta_0 X$,

$$Z[O_c^p] = \{Z' \cap X : Z' \in Z_c[\beta_0 X], p \in int_{\beta_0 X} Z'\}.$$

Proof. Suppose that $Z \in Z[O_c^p]$, i.e., $p \in int_{\beta_0 X} cl_{\beta_0 X} Z$. By Corollary 1.4, there exists $Z' \in Z_c[\beta_0 X]$ such that $Z' \cap X = Z$. Clearly $p \in int_{\beta_0 X} Z'$. Hence Z belongs to the right hand side set. Now let $Z' \in Z_c[\beta_0 X]$ and $p \in int_{\beta_0 X} Z'$. There exists a clopen set $U \subseteq \beta_0 X$ such that $p \in U \subseteq int_{\beta_0 X} Z'$. Therefore $U \cap X \subseteq Z' \cap X$ and hence $p \in int_{\beta_0 X} cl_{\beta_0 X}(Z' \cap X)$. This implies that $Z' \cap X \in Z[O_c^p]$. □

We denote by $C_c^K(X)$, the set of all functions of $C_c(X)$ with compact support. In the following we characterize the set $C_c^K(X)$ as an ideal of $C_c(X)$.

Theorem 2.4. For a zero-dimensional space X, $C_c^K(X) = O_c^{\beta_0 X \setminus X}$.

Proof. Suppose that $f \in C_c^K(X)$. Then $cl_X (X \setminus Z(f))$ is compact and hence $cl_{\beta_0 X} (X \setminus Z(f)) \subseteq X$. By Corollary 1.4, there exists $Z' \in Z_c[\beta_0 X]$ such that $Z(f) = Z' \cap X$. Therefore $X \cap (\beta_0 X \setminus Z') = X \setminus Z(f)$. Since X is dense and $\beta_0 X \setminus Z'$ is open in $\beta_0 X$, $cl_{\beta_0 X} (\beta_0 X \setminus Z') \subseteq X$. Thus $\beta_0 X \setminus X \subseteq \beta_0 X \setminus cl_{\beta_0 X} (\beta_0 X \setminus Z') = int_{\beta_0 X} Z'$. Hence by Proposition 2.3, for all $p \in \beta_0 X \setminus X$, $Z' \cap X \in Z_c[O_c^p]$ and therefore $f \in O_c^{\beta_0 X \setminus X}$. For the reverse inclusion, suppose that $f \in O_c^{\beta_0 X \setminus X}$. Then $\beta_0 X \setminus X \subseteq int_{\beta_o X} cl_{\beta_0 X} Z(f)$. By Proposition 2.3, there exists $Z' \in Z_c[\beta_0 X]$ such that $Z' \cap X = Z(f)$ and $\beta_0 X \setminus X \subseteq int_{\beta_0 X} Z'$. This implies that $\beta_0 X \setminus int_{\beta_0 X} Z' \subseteq X$. Since $\beta_0 X \setminus int_{\beta_0 X} Z' = cl_{\beta_0 X}(\beta_0 X \setminus Z') \subseteq X$ and $X \setminus Z(f) \subseteq \beta_0 X \setminus Z'$, clearly $cl_X(X \setminus Z(f))$ is compact and hence $f \in C_c^K(X)$. □

The following two results are important for the rest of this section. We recall that a subset $S \subseteq X$ is C_c-embedded in X if for each $f \in C_c(S)$, there exists $F \in C_c(X)$ such that $F|_X = f$.

Proposition 2.5. If $S \subseteq X$ is C_c-embedded in X, then it is separated by a clopen partition from every $Z \in Z_c[X]$ disjoint from it.

Proof. Suppose that $h \in C_c(X)$ and $S \cap Z(h) = \emptyset$. Define $f(s) = \frac{1}{h(s)}$ for all $s \in S$. Let $F \in C_c(X)$ be such that $F|_S = f$. Put $k = hF$. Clearly $k \in C_c(X)$ and $k|_S = 1$ and $k|_{Z(h)} = 0$. It is enough to choose $0 < r < 1$ such that $r \notin k(X)$. Then $U = k^{-1}(-\infty, r)$ is a clopen subset of X that $Z(h) \subseteq U$ and $S \subseteq X \setminus U$. □

In the following we need to know about the existence of C_c-embedded subsets. Note that C-embedded subsets are not applicable for our purpose. Because we do not know anything about the cardinality of the image of the extension of a function with countable image. The following lemma gives us a condition for existing a C_c-subset in a zero-dimensional space.

Lemma 2.6. Let X be zero-dimensional and $f \in C_c(X)$ carries a subset $S \subseteq X$ homeomorphically to a closed subset $f(S) \subseteq f(X)$. Then S is C_c-embedded in X.

Proof. Suppose that $f|_S : S \to f(S)$ is a homeomorphism. Therefore $f^{-1} : f(S) \to S$ is continuous. Consider $g \in C_c(S)$. We observe that the composition gof^{-1} belongs to $C_c(f(S))$. The subset $f(X)$ is countable and therefore normal. Hence gof^{-1} has an extension G to all of $f(X)$. It is obvious to see that $G \in C_c(f(X))$. The composition Gof belongs to $C_c(X)$. For $s \in S$, we have $Gof(s) = G(f(s)) = gof^{-1}(f(s)) = g(s)$. Therefore Gof is the extension of g to X which has countable image. □

The previous lemma leads us to the following corollary.

Corollary 2.7. For a zero-dimensional space X, let $E \subseteq X$. Suppose the function $h \in C_c(X)$ is unbounded on E. Then E contains a copy of $\mathbb{N}$, which is C_c-embedded in X and h approach to infinity on E.

In the following, we adapt the original approach of Mandelker in [11] and [12] to functionally countabe subalgebras for characterizing the subset consisting of all functions in $C_c(X)$ with pseudocompact support. We denote by $C_c^{\psi}(X)$, the set of all functions in $C_c(X)$ with pseudocompact support. We recall that a subset $S \subseteq X$ is relatively pseudocompact with respect to $C_c(X)$, if for each $f \in C_c(X)$, the function $f|_S$ is bounded. We have the following equivalence for relatively pseudocompact subsets with respect to $C_c(X)$.

Proposition 2.8. Let X be a zero-dimensional space. The set $A \subseteq X$ is relatively pseudocompact with respect to $C_c(X)$ if and only if $cl_{\beta_0 X} A \subseteq \upsilon_0 X$.

Proof. For the necessity, suppose $cl_{\beta_0 X} A \cap (\beta_0 X \setminus \upsilon_0 X) \neq \emptyset$ and choose $p \in cl_{\beta_0 X} A \cap (\beta_0 X \setminus \upsilon_0 X)$. There exists a sequence $\{V_n : n \in \mathbb{N}\}$ of clopen neighborhoods of p in $\beta_0 X$ such that $(\cap_{n=1}^{\infty} V_n) \cap \upsilon_0 X = \emptyset$. By part (a) of Lemma 1.3, there exists $F \in C_c(\beta_0 X)$ such that $Z(F) = \cap_{n=1}^{\infty} V_n$. Put $f = F|_X$ and define $h = \frac{1}{f}$. Clearly $h \in C_c(X)$ and since p is a limit point of A in $\beta_0 X$, h is unbounded on A, which is a contradiction. For the sufficiency, consider $f \in C_c(X)$. The extension $f^{\upsilon_0} \in C_c(\upsilon_0 X)$ is bounded on the compact subset $cl_{\beta_0 X} A$. Therefore f should be bounded on A. □

For continuing our progress, we need a proposition which is due to Pierce. The reader could find it in [18, Lemma 1.9.3].

Proposition 2.9. A zero-dimensional space X is not pseudocompact if and only if there exists a continuous and onto map $f : X \to \mathbb{N}$.

Theorem 2.10. If $f \in C_c(X)$ and $X \setminus Z(f)$ is relatively pseudocompact with respect to $C_c(X)$, then $X \setminus Z(f)$ is pseudocompact.

Proof. Suppose on the contrary that $S = cl_X(X \setminus Z(f))$ is not pseudocompact. Hence by Proposition 2.9, there exists $h \geq 1$ and $h \in C_c(S)$ that is unbounded on $X \setminus Z(f)$. By Corollary 2.7, there exists a countable discrete subset $D \subseteq X \setminus Z(f)$ such that D is C_c-embedded in S and h is unbounded on D. Proposition 2.5 implies that there exists a clopen set $O \subseteq S$ such that $D \subseteq O$ and $Z(f) \cap S \subseteq S \setminus O$. Define the function k as follows,

$$k(x) = \begin{cases} \frac{1}{h(s)} \vee \chi_{S \setminus O}(s), & s \in S \\ 1, & s \in cl_X\left(X \setminus S\right). \end{cases}$$

Pasting lemma implies that k is continuous on X. Clearly $k \in C_c(X)$ and $k > 0$. Hence $\frac{1}{k}$ is unbounded on D. Thus S is not relatively pseudocompact with respect to $C_c(X)$, a contradiction. □

Theorem 2.11. Let X be a zero-dimensional space and $f \in C_c(X)$. If $\beta_0 X \setminus \upsilon_0 X \subseteq cl_{\beta_0 X} Z(f)$, then $\beta_0 X \setminus \upsilon_0 X \subseteq int_{\beta_0 X} cl_{\beta_0 X} Z(f)$

Proof. Suppose that $p \in \beta_0 X \setminus \upsilon_0 X$. There exists a sequence $\{V_n : n \in \mathbb{N}\}$ of clopen neighborhoods of p in $\beta_0 X$ such that $(\cap_{n=1}^{\infty} V_n) \cap \upsilon_0 X = \emptyset$. By part (a) of Lemma 1.3, there exists $h \in C_c(\beta_0 X)$ such that $Z(h) = \cap_{n=1}^{\infty} V_n$. Put $T = \beta_0 X \setminus Z(h)$. Consider the function $\frac{1}{h}$ on T. If $Z(h)$ meets $cl_{\beta_0 X}(X \setminus Z(f))$, then $\frac{1}{h}$ which is continuous on T is unbounded on $X \setminus Z(f)$. Hence by Corollary 2.7, there exists a countable closed set $S \subseteq X \setminus Z(f)$ which is C_c-embedded in T. Thus S is C_c-embedded in X and by Proposition 2.5, there exists a clopen set $O \subseteq X$ such that $S \subseteq O$ and $Z(f) \subseteq X \setminus O$. Therefore $cl_{\beta_0 X} S \cap cl_{\beta_0 X} Z(f) = \emptyset$. Since S is closed in T and also is noncompact, there exists $p \in cl_{\beta_0 X} S \setminus T$. This implies that $p \in Z(h)$ and $p \notin cl_{\beta_0 X} Z(f)$ which is a contradiction. Hence $Z(h) \cap cl_{\beta_0 X}(X \setminus Z(f)) = \emptyset$ and so $Z(h) \subseteq \beta_0 X \setminus cl_{\beta_0 X}(X \setminus Z(f)) \subseteq cl_{\beta_0 X} Z(f)$. Therefore $cl_{\beta_0 X} Z(f)$ is a neighborhood of p. □

Now we are ready to characterize another important ideal in $C_c(X)$. We denote by $C_c^{\psi}(X)$, the set of all functions in $C_c(X)$ with pseudocompact support.

Theorem 2.12. Let X be a zero-dimensional space. Then $C_c^{\psi}(X) = M_c^{\beta_0 X \setminus \upsilon_0 X} = O_c^{\beta_0 X \setminus \upsilon_0 X}$.

Proof. By Theorem 2.11, the second equality is clear. Assume that $f \in C_c^{\psi}(X)$. Since $S(f) = X \setminus Z(f)$ is pseudocompact and $(\upsilon_0 X \setminus Z(f^{\upsilon_0})) \cap X = X \setminus Z(f)$, the subset $\upsilon_0 X \setminus Z(f^{\upsilon_0})$ is also pseudocompact. Note that $S(f^{\upsilon_0}) = \upsilon_0 X \setminus Z(f^{\upsilon_0})$ is a cozero-set of the real compact space $\upsilon_0 X$, and hence it should be realcompact, see [6, Corollary 8.14]. Therefore $S(f^{\upsilon_0})$ must be compact. We observe that

$$\beta_0 X = cl_{\beta_0 X} Z(f) \cup cl_{\beta_0 X} S(f) = cl_{\beta_0 X} Z(f) \cup S(f^{\upsilon_0}),$$

and hence we have $\beta_0 X \setminus \upsilon_0 X \subseteq cl_{\beta_0 X} Z(f)$. Thus $f \in M_c^{\beta_0 X \setminus \upsilon_0 X}$. Now if $f \in O_c^{\beta_0 X \setminus \upsilon_0 X}$, there exists a compact set K such that

$$\beta_0 X \setminus \upsilon_0 X \subseteq \beta_0 X \setminus K \subseteq cl_{\beta_0 X} Z(f),$$

and hence $X \setminus Z(f) \subseteq \beta_0 X \setminus cl_{\beta_0 X} Z(f) \subseteq K \subseteq \upsilon_0 X$. Since $X \setminus Z(f)$ is relatively pseudocompact with respect to $C_c(X)$, by Theorem 2.10, the set $X \setminus Z(f)$ is pseudocompact. □

For an $\mathbb{N}$-compact space, Theorem 2.4 and Theorem 2.12 imply the following corollary which says that in an $\mathbb{N}$-compact space, $C_c^K(X)$ equals to that intersection of all free maximal ideals in $C_c(X)$.

Corollary 2.13. If X is an $\mathbb{N}$-compact space, then $C_c^K(X) = M_c^{\beta_0 X \setminus X}$.

3. REMAINDER OF BANASCHEWSKI COMPACTIFICATION VIA $C_c(X)$.

In this section, first we want to find the least cardinality of the remainder $\beta_0 X \setminus \upsilon_0 X$. In other point of view, we show that for a zero-dimensional non pseudocompact space X, we have at least $2^{2^{\aleph_0}}$ clopen ultrafilters on X which do not have countable intersection property. In the rest of this section we observe that the remainder of the Banaschewski compactification of an $\mathbb{N}$-compact space is an almost P-space and a connection between the remainder and Parovičenko spaces are given. Finally, for a zero-dimensional, locally compact space X which is not pseudocompact, we give a lower bound for the cellularity of spaces $\beta_0 X \setminus \upsilon_0 X$ and $\beta_0 X \setminus X$. The following two lemmas are needed in the sequel. The second one is a consequence of Proposition 2.9.

Lemma 3.1. Each C_c-embedded subset S of X is two-embedded.

Proof. Let $f : S \to \{0,1\}$ be a continuous two-valued function, then $f \in C_c(X)$. Therefore there exists $G \in C_c(X)$ such that $G|_S = f$. The image of G is countable and there exists a real number $0 < r < 1$ such that $r \notin G(X)$. The subset $U = G^{-1}(\infty, r)$ is clopen in X and $f^{-1}(0) \subseteq U$ and $f^{-1}(1) \subseteq X \setminus U$. Define $F : X \to \{0,1\}$ to be 0 on U and 1 on $X \setminus U$. The function F is continuous and two-valued whose restriction to S is f. □

Lemma 3.2. Let X be a zero-dimensional space. Then X is pseudocompact if and only if $\beta_0 X = \upsilon_0 X$.

Proof. For the necessity, suppose that X is pseudocompact and $\beta_0 X \setminus \upsilon_0 X \neq \emptyset$. For $p \in \beta_0 X \setminus \upsilon_0 X$, there exists a countable set consisting of clopen neighborhoods of p, say $\{U_n : n \in \mathbb{N}\}$, in $\beta_0 X$ with $(\cap_{n=1}^{\infty} U_n) \cap \upsilon_0 X = \emptyset$. By part (a) of Lemma 1.3, there exists a function $f \in C_c(\beta_0 X)$ such that $Z(f) = \cap_{n=1}^{\infty} U_n$. If we restrict f to X, then for the function $g = f|_X \in C_c(X)$, we have $Z(g) = \emptyset$. Hence g is a unit of $C_c(X)$. The function $\frac{1}{g}$ belongs to $C_c(X)$ and clearly g is unbounded on X

which is a contradiction.
For the sufficiency, suppose that X is not pseudocompact. Hence by Proposition 2.9, there exists a continuous and onto map $f : X \to \mathbb{N}$. The function f has countable image and therefore has an extension to $\upsilon_0 X$. But $\upsilon_0 X$ is compact and the extension of f is unbounded which is a contradiction. This completes the proof of our lemma. □

Now we are ready to present one of our main theorems in this section. This result is notable whenever we deal with zero-dimensional spaces which are not strongly zero-dimensional.

Theorem 3.3. For a zero-dimensional space X, let $f \in C_c(\beta_0 X)$ and $Z(f) \cap X = \emptyset$. Then $Z(f)$ contains a copy of $\beta\mathbb{N}$, the Stone-Čech compactification of $\mathbb{N}$, and therefore its cardinality is at least $2^{2^{\aleph_0}}$.

Proof. Denote the restriction of f to X by g. So $Z(g) = \emptyset$ and g belongs to $C_c(X)$. This function has an inverse in $C_c(X)$, say $\frac{1}{g}$. Obviously $\frac{1}{g}$ is unbounded on X. Hence by Corollary 2.7, there exists a C_c-embedded copy of $\mathbb{N}$ in X on which the function $\frac{1}{g}$ tends to infinity. Since $\mathbb{N}$ is C_c-embedded in X, by Lemma 3.1, it is also two-embedded in X and therefore two-embedded in $\beta_0 X$. Thus every continuous two-valued function on $\mathbb{N}$ is two-embedded in $cl_{\beta_0 X}\mathbb{N}$. By the uniqueness theorem in the existence of Banaschewski compactification, $cl_{\beta_0 X}\mathbb{N}$ and $\beta_0\mathbb{N}$ are homeomorphic, and also we have $cl_{\beta_0 X}\mathbb{N} \setminus \mathbb{N} \subseteq Z(f)$. The discrete space $\mathbb{N}$ is strongly zero-dimensional and hence $\beta_0\mathbb{N} = cl_{\beta_0 X}\mathbb{N} = \beta\mathbb{N}$. The cardinality of $\beta\mathbb{N}$ equals to $2^{2^{\aleph_0}}$ and therefore $|Z(f)| \geq 2^{2^{\aleph_0}}$. □

Since each G_δ-point is a zero-set, Lemma 1.3 and Theorem 3.3 imply the following corollary.

Corollary 3.4. For a zero-dimensional space X, no point $p \in \beta_0 X \setminus X$ is a G_δ-point of $\beta_0 X$.

In the next result, by applying Theorem 3.3, we give the least cardinality of the remainder $\beta_0 X \setminus \upsilon_0 X$, whenever X is zero-dimensional and non pseudocompact space.

Proposition 3.5. Let X be a zero-dimensional and non pseudocompact space. The remainder $\beta_0 X \setminus \upsilon_0 X$ has at least $2^{2^{\aleph_0}}$ points.

Proof. Since X is a zero-dimensional and non pseudocompact space, By Proposition 2.9, there exists a continuous and onto function $f : \upsilon_0 X \to \mathbb{N}$. Therefore the function $g = \frac{1}{f}$ is continuous whose image equals to the set $\{\frac{1}{n} : n \in \mathbb{N}\}$. Note that the set $\{\frac{1}{n} : n \in \mathbb{N}\} \cup \{0\}$ is zero-dimensional and compact. Hence by [19, 4.7, proposition (d)], g has a continuous extension $G : \beta_0 X \to \{\frac{1}{n} : n \in \mathbb{N}\} \cup \{0\}$. For each $n \in \mathbb{N}$, choose x_n such that $g(x_n) = \frac{1}{f(x_n)} = \frac{1}{n}$. Each $g^{-1}(\frac{1}{n})$ is clopen in $\upsilon_0 X$ and hence the cluster points of the set $\{x_n : n \in \mathbb{N}\}$ is contained in $\beta_0 X \setminus \upsilon_0 X$. If we consider a cluster point p of the set $\{x_n : n \in \mathbb{N}\}$, then we observe that $G(p) = 0$. Therefore $G \in C_c(\beta_0 X)$ and $Z(G) \neq \emptyset$. Since $Z(G) \cap \upsilon_0 X = \emptyset$, Theorem 3.3 implies that $|Z(G)| \geq 2^{2^{\aleph_0}}$. Thus the cardinality of $\beta_0 X \setminus \upsilon_0 X$ is at least $2^{2^{\aleph_0}}$. □

We recall that the local weight of the space X at a point p, denoted $\chi(p, X)$ or $\chi(p)$, is the least cardinal equal to the cardinal number of a (filter) base for the neighborhoods of p. By Proposition 3.5, for a zero-dimensional non pseudocompact space X, we have the following corollary.

Corollary 3.6. Let X be a zero-dimensional and non pseudocompact space. Then;
(a) If $p \in \beta_0 X \setminus \upsilon_0 X$, then $\{p\}$ is not a zero-set of $\beta_0 X \setminus \upsilon_0 X$;
(b) $\beta_0 X \setminus \upsilon_0 X$ has no isolated point;
(c) The local weight of $\beta_0 X \setminus \upsilon_0 X$ at each of its points is uncountable.

Proof. (a) If p is a zero-set of $\beta_0 X \setminus \upsilon_0 X$, there exists a sequence $\{U_n : n \in \mathbb{N}\}$ of clopen neighborhoods of p in $\beta_0 X$ such that $\{p\} = (\cap_{n=1}^{\infty} U_n) \cap (\beta_0 X \setminus \upsilon_0 X)$. Also there exists a sequence $\{V_n : n \in \mathbb{N}\}$ of clopen neighborhoods of p in $\beta_0 X$ such that $(\cap_{n=1}^{\infty} V_n) \cap \upsilon_0 X = \emptyset$. Hence $\{p\} = (\cap_{n=1}^{\infty} U_n) \cap (\cap_{n=1}^{\infty} V_n)$. Part (a) of Lemma 1.3 implies that there exists $f \in C_c(\beta_0 X)$ such that $\{p\} = Z(f)$. But this contradicts with Proposition 3.5.
(b) Since each isolated point is a zero-set, clearly part (a) implies part (b).
(c) If for some p the locall weight at p is countable, then p is a G_δ-point and hence a zero-set of $\beta_0 X \setminus \upsilon_0 X$, which contradicts with part (a). □

We recall that a topological space is scattered if each of its nonempty subsets has an isolated point with the relative topology. Lemma 3.2 together with part (b) of Corollary 3.6, imply the following corollary.

Corollary 3.7. For a zero-dimensional space X, if $\beta_0 X$ is scattered then X is pseudocompact.

In the following, we observe some results about connection between Banaschewski compactification and almost P-spaces. We recall that a topological space X is an almost P-space if each nonempty G_δ-subset of X has nonempty interior, see [10] and [20].

Proposition 3.8. $\beta_0 X$ is an almost P-space if and only if X is pseudocompact and an almost P-space.

Proof. For the necessity, suppose that $\beta_0 X$ is an almost P-space and X is not pseudocompact. For $p \in \beta_0 X \setminus \upsilon_0 X$, there exist a sequence of countable clopen subsets of X, say $\{U_n : n \in \mathbb{N}\}$, such that $\cap_{n=1}^{\infty} U_n = \emptyset$ and for each $n \in \mathbb{N}$, $p \in cl_{\beta_0 X} U_n$. Note that for each $n \in \mathbb{N}$, $cl_{\beta_0 X} U_n$ is a clopen subset of $\beta_0 X$. Hence $G = \cap_{n=1}^{\infty} cl_{\beta_0 X} U_n$ is a G_δ-subset in $\beta_0 X$. By our hypothesis, $int_{\beta_0 X} G \neq \emptyset$. Therefore G must intersects X. But $G \cap X = \cap_{n=1}^{\infty} U_n = \emptyset$, which is a contradiction. Now we show that X is an almost P-space. For a nonempty G_δ-set G, there exists a sequence $\{V_n : n \in \mathbb{N}\}$ of open subsets of X such that $G = \cap_{n=1}^{\infty} V_n$. For $p \in G$, there exists a sequence of clopen sets $\{O_n : n \in \mathbb{N}\}$ such that for each $n \in \mathbb{N}$, $p \in O_n \subseteq V_n$. For each $n \in \mathbb{N}$, $cl_{\beta_0 X} O_n$ is clopen in $\beta_0 X$ and $p \in T = \bigcap_{n=1}^{\infty} (cl_{\beta_0 X} O_n)$. So there exists a nonempty open set V in $\beta_0 X$ such that $V \subseteq T$. The set $V \cap X$ is nonempty and is contained entirely in $\cap_{n=1}^{\infty} O_n \subseteq G$. Therefore X is an almost P-space.
For the sufficiency, let H be a nonempty G_δ-subset of $\beta_0 X$. There exist a sequence $\{V_n : n \in \mathbb{N}\}$ of clopen subsets of $\beta_0 X$ and $p \in H$ such that $p \in \cap_{n=1}^{\infty} V_n \subseteq H$. For each $n \in \mathbb{N}$, $V_n \cap X$ is clopen in X and the collection $\Omega = \{V_n \cap X : n \in \mathbb{N}\}$ has finite intersection property. Since $\beta_0 X = \upsilon_0 X$, Ω is contained in a clopen ultrafilter with countable intersection property. Hence $\bigcap_{n=1}^{\infty} (V_n \cap X)$ is nonempty and therefore a G_δ-subset of X. So there exists a neighborhood O in $\beta_0 X$ such that $O \cap X \subseteq \bigcap_{n=1}^{\infty} (V_n \cap X)$. By taking closure, we observe that $O \subseteq \cap_{n=1}^{\infty} V_n$. Hence H has nonempty interior. □

In the following theorem, we introduce a class of almost P-spaces in connection with Banaschewski compactification of locally compact and $\mathbb{N}$-compact spaces.

Theorem 3.9. Let X be a locally compact and $\mathbb{N}$-compact space. Then $\beta_0X \setminus X$ is an almost P-space.

Proof. Let G be a nonempty G_δ-set in $\beta_0X \setminus X$ and $p \in G$. There exists a sequence of open sets in β_0X, say $\{V_n : n \in \mathbb{N}\}$, such that $G = (\bigcap_{n=1}^{\infty} V_n) \cap (\beta_0X \setminus X)$. Also there exists a sequence of clopen sets $\{U_n : n \in \mathbb{N}\}$ of X such that $\bigcap_{n=1}^{\infty} U_n = \emptyset$ and $p \in \bigcap_{n=1}^{\infty} cl_{\beta_0X}U_n$. Note that the G_δ-set $\bigcap_{n=1}^{\infty} cl_{\beta_0X}U_n$ is a subset of $\beta_0X \setminus X$. Obviously $H = (\bigcap_{n=1}^{\infty} cl_{\beta_0X}U_n) \cap (\bigcap_{n=1}^{\infty} V_n)$ is a G_δ-subset of β_0X and $p \in H \subseteq \beta_0X \setminus X$. One could find a sequence of clopen subsets of β_0X, say $\{O_n : n \in \mathbb{N}\}$, such that $p \in \bigcap_{n=1}^{\infty} O_n \subseteq H$. By part (a) of Lemma 1.3, there exists $F \in C_c(\beta_0X)$ such that $Z(F) = \bigcap_{n=1}^{\infty} O_n$. Since X is locally compact, for each $i \in \mathbb{N}$, there exists an open set $W_i \subseteq X$ such that cl_XW_i is compact and for each $x \in W_i$, $F(x) \leq \frac{1}{i}$. It is enough to show that the set $(\beta_0X \setminus X) \cap cl_{\beta_0X}\left(\bigcup_{i\in\mathbb{N}} W_i\right)$ is a subset of $Z(F)$. Consider $t \in (\beta_0X \setminus X) \cap cl_{\beta_0X}\left(\bigcup_{i\in\mathbb{N}} W_i\right)$. Each neighborhood P of t in β_0X intersects infinitely many W_i's. To see this, assume that for a neighborhood P of t, there exists $n \in \mathbb{N}$ such that for all $m > n$; $P \cap W_m = \emptyset$. This implies that $t \in cl_{\beta_0X}\left(\bigcup_{i=1}^{n} W_i\right)$. Each W_i has compact closure in X, and hence $cl_{\beta_0X}\left(\bigcup_{i=1}^{n} W_i\right)$ is a compact subset of X which implies that $t \in X$ and this is a contradiction with the choice of t. Since each neighborhood P of t in β_0X intersects infinitely many W_i's, the function F must vanishes in t. Hence

$$T = (\beta_0X \setminus X) \cap (cl_{\beta_0X}(\cup_{i\in\mathbb{N}}W_i)) \subseteq Z(F).$$

It is easy to see that T is nonempty, open in $\beta_0X \setminus X$ and $T \subseteq G$, which shows that G has nonempty interior. This completes the proof. □

Using Corollary 3.6 together with Theorem 3.9, we could find a class of spaces whose appearance is in Boolean algebras, see for example [4]. We recall that a compact zero-dimensional space X is called a Parovičenko space if it has the following properties:
(a) X has no isolated points.
(b) Nonempty G_δ-sets have nonempty interiors.
(c) X is an F-space (i.e., for each $f \in C(X)$, the subsets $pos(f)$ and $neg(f)$ are completely separated).

The following proposition gives us a large class of Parovičenko spaces in Banaschewski compactification of a non pseudocompact zero-dimensional space.

Proposition 3.10. Let Z be a zero-set of β_0X such that $Z \cap X = \emptyset$. Then Z is a Parovičenko space.

Proof. Evidently Z is compact and zero-dimensional. Since Z is a G_δ-subset of β_0X, $W = \beta_0X \setminus Z$ is σ-compact and therefore strongly zero-dinensional, see [6, 16.17]. Thus $\beta_0W = \beta W$. The space W is Lindelöf and hence is $\mathbb{N}$-compact. We could observe that $Z = \beta_0W \setminus W = \beta W \setminus W$. So by Theorem 3.9, Z must be an almost P-space. Also since W is locally compact and σ-compact, by Theorem 14.27 of [6], Z is an F-space. Part (b) of Corollary 3.6, implies that Z contains no isolated point. Hence Z must be a Parovičenko space. □

Let X be zero-dimensional, locally compact space which is not pseudocompact. We close this section by giving a lower bound for the cellularity of the subspaces

$\beta_0 X \setminus \upsilon_0 X$ and $\beta_0 X \setminus X$. We recall that the cellularity of a space Y, denoted by $c(Y)$, is the smallest cardinal number κ for which each pairwise disjoint family of nonempty open sets of Y has κ or fewer members. Also we remind that for a zero-dimensional space X and a cardinal number κ, a partition of X with length κ is a family $\{U_i : i \in \mathbb{I}\}$ of pairwise disjoint clopen subsets of X whose union is X and $|\mathbb{I}| = \kappa$. Note that by Proposition 2.9, every zero-dimensional space which is not pseudocompact, has a partition with length $\aleph_0$. The following theorem which is due to Tarski, is needed for our purpose, see [8].

Theorem 3.11. [Tarski] Let E be an infinite set. Then there is a collection $\Re$ of subsets of E such that $|\Re| = |E|^{\aleph_0}$, $|R| = \aleph_0$ for each $R \in \Re$ and the intersection of any two distinct members of $\Re$ is finite.

Theorem 3.12. Let X be a zero-dimensional and locally compact space which is not pseudocompact. If X has a partition of length κ, then the cellularity of subspaces $\beta_0 X \setminus \upsilon_0 X$ and $\beta_0 X \setminus X$ are at least $\kappa^{\aleph_0}$.

Proof. We just prove that the cellularity of $\beta_0 X \setminus \upsilon_0 X$ is at least $\kappa^{\aleph_0}$. The second one can be derived similarly. Let $\{U_i : i \in \mathbb{I}\}$ be a partition of X with $|\mathbb{I}| = \kappa$. For each $i \in \mathbb{I}$, choose a nonempty clopen and compact subset $W_i \subseteq U_i$. Note that $\{cl_{\upsilon_0 X} U_i : i \in \mathbb{I}\}$ is a partition of length κ for $\upsilon_0 X$ and for each $i \in \mathbb{I}$, $cl_{\upsilon_0 X} W_i = W_i$ is clopen in $\upsilon_0 X$. For each subset $J \subseteq \mathbb{I}$, define

$$A(J) = (\beta_0 X \setminus \upsilon_0 X) \cap \left(cl_{\beta_0 X}(\bigcup_{i \in J} W_i) \right).$$

We remind that since for each $J \subseteq \mathbb{I}$, $\{W_i : i \in J\}$ is a locally finite family of clopen subsets of $\upsilon_0 X$, then $\bigcup_{i \in J} W_i$ is clopen in $\upsilon_0 X$ (see [2, Theorem 1.1.11]) and hence $cl_{\beta_0 X}(\bigcup_{i \in J} W_i)$ is clopen in $\beta_0 X$. This implies that for each subset $J \subseteq \mathbb{I}$, $A(J)$ is clopen in $\beta_0 X \setminus \upsilon_0 X$. It is clear that for each two subsets J_1, J_2 of $\mathbb{I}$, $A(J_1) \cap A(J_2) = A(J_1 \cap J_2)$. For a subset $J \subseteq \mathbb{I}$, we claim that $A(J) = \emptyset$ if and only if J is finite. For, if $J \subseteq \mathbb{I}$ is finite, then $cl_{\beta_0 X}(\bigcup_{i \in J} W_i) = \bigcup_{i \in J} W_i$ and hence $A(J) = \emptyset$. For the converse, if $J \subseteq \mathbb{I}$ is infinite, then for each $j \in J$, choose $x_j \in W_j$. Clearly the subset $B = \{x_j : j \in J\}$ is a closed and discrete subset of $\upsilon_0 X$. Since $\beta_0 X$ is compact, the subset B has a cluster point p in $\beta_0 X \setminus \upsilon_0 X$. This implies that $A(J) \neq \emptyset$.
Now apply Theorem 3.12 to find a collection Υ of $\kappa^{\aleph_0}$ infinite subsets of $\mathbb{I}$ such that any two members of Υ have finite intersection. Define $\mathcal{T} = \{A(J) : J \in \Upsilon\}$. Evidently $\mathcal{T}$ contains $\kappa^{\aleph_0}$ pairwise disjoint clopen subsets of $\beta_0 X \setminus \upsilon_0 X$. This implies that $c(\beta_0 X \setminus \upsilon_0 X) \geq \kappa^{\aleph_0}$. □

By applying Proposition 2.9 and Theorem 3.12, the following corollary is immediate.

Corollary 3.13. Let X be a zero-dimensional, locally compact space which is not pseudocompact. Then the cellularity of spaces $\beta_0 X \setminus \upsilon_0 X$ and $\beta_0 X \setminus X$ are at least $2^{\aleph_0}$.

REFERENCES

[1] P. Bhattacharjee, M. L. Knox, W. W. McGovern, *The classical ring of quotients of $C_c(X)$* , Appl. Gen. Topol. 15, no. 2(2014), 147-154

[2] R. Engelking , *General topology*, PWN-Polish Sci. Publ., 1977.

[3] R. Engelking, S. Mrowka, *On E-compact spaces* , Bull. Acad. Polon. Sci, 6 (1958), 429-436.

[4] R. Frankiewicz, P. Zbierski, *Hausdorff gaps and limits*, Studies in logic and the foundations of mathematics, 132, North- Holland, (1994).

[5] M. Ghadermazi, O. A. S. Karamzadeh, M. Namdari, *On the functionally countable subalgebra of* $C(X)$, Rend. Sem. Mat. Univ. Padova, 129 (2013), 47-69.

[6] L. Gillman, M. Jerison, *Rings of Continuous Functions*, Springer-Verlag, 1976.

[7] Hager A., Kimber C., McGovern W. W., *Unique a-closure for some l-groups of rational valued functions* , Czech. Math. J. 55 (2005), 409-421.

[8] R. E. Hodel, Jr., *Cardinal functions I*, in: K. Kunen and J. E. Vaughan (eds.), Hand- book of Set-Theoretic Topology, North-Holland, Amsterdam, (1984), 1-61.

[9] D. G. Johenson, M. Mandelker, *Functions with pseudocompact support* , General topology appl, 3(1973) 331-338

[10] R. Levy, *Almost P-spaces* , Canad. J. Math. 2 (1977), 284-288.

[11] M. Mandelker, *Round z-filters and round subsets of* βX , Israel. J. Math. 7 (1969), 1-8.

[12] M. Mandelker, *Supports of continuous functions* , Trans. Amer. Math. Soc. 156 (1971), 73-83.

[13] S. Mrowka, *Structures of continuous functions. III, Real and lattices of integer-valued continuous functions* , Vehr. Nederl. Akad. Weten., Sect. I, 68 (1965), 74-82.

[14] S. Mrowka, S. D. Shore, *Structures of continuous functions. V, On homomorphisms of structures of continuous functions with zero-dimensional compact domain* , Vehr. Nederl. Akad. Weten., Sect. I, 68 (1965), 92-94.

[15] S. Mrowka, *On E-compact spaces. II* , Bull. Acad. Polon. Sci, 14 (1966), 597-605.

[16] S. Mrowka, *Further results on E-compact spaces. I* , Acta. Math. Hung. 120 (1968), 161-185.

[17] S. Mrowka, *Structures of continuous functions. I* , Acta. Math. Hung. 21(3-4) (1970), 239-259.

[18] R. S. Pierce, *Rings of integer-valued continuous functions* , Trans. Amer. Math. Soc., 100 (1961), 371-394.

[19] J. R. Porter, R. G. Woods, *Extensions and Absolutes of Hausdorff spaces*, Springer-Verlag, 1988.

[20] A. I. Veksler, *P'-points, P'-sets, P'-spaces. A new class of order-continuous measures and functionals* , Sov. Math. Dokl. 14 (1973), 1445-1450.

Department of Mathematics, Yasouj University, Yasouj, Iran
E-mail address: `alireza.olfati@yu.ac.ir, olfati.alireza@gmail.com`